

Acoustic Complexity Index to assess Benthic Biodiversity of a Partially Protected Area in the Southwest of the UK.

Bede F. R. **Davies**^a, Martin J. Attrill^a, Luke Holmes^a, Adam Rees^a, Matthew J. Witt^b and
Emma V. Sheehan^a.

^a University of Plymouth, Drake Circus, Plymouth, United Kingdom, PL4 8AA

^b University of Exeter, Hatherly Laboratories, College of Life & Environmental Sciences,
Prince of Wales Road, Exeter, United Kingdom, EX4 4PS

Correspondence author: Bede Ffinian Rowe Davies, bede.davies@plymouth.ac.uk

Highlights:

- Acoustic Complexity Index higher in fished area vs protected area
- Acoustic Complexity Index does not correlate with simultaneous biodiversity indices
- Acoustic Complexity Index covaries with mobile assemblage composition

Abstract

The soundscape of the marine environment is a relatively understudied area of ecology that has the potential to provide large amounts of information on biodiversity, reproductive behaviour, habitat selection, spawning and predator-prey interactions. Biodiversity is often visually assessed

and used as a proxy for ecosystem health. Visual assessment using divers or remote video methods can be expensive, and limited to times of good weather and water visibility. Previous studies have concluded that acoustic measures, such as the Acoustic Complexity Index (ACI), correlate with visual biodiversity estimates and offer an alternative to assess ecosystem health.

Here, the ACI measured over 5 years in a Marine Protected Area (MPA) in the UK, Lyme Bay, was analysed alongside another monitoring method, Baited Remote Underwater Video Systems (BRUVs). Two treatments were sampled annually in the summer from 2014 until 2018 with sites inside the MPA, as well as Open Control sites outside of the MPA.

Year by year correlations, which have been used elsewhere to test ACI, showed significant correlations with Number of Species and ACI. However, the sign of these correlations changed almost yearly, showing that more in-depth analyses are needed.

Multivariate analysis of the benthic assemblage composition (from BRUVs) was carried out by Permutational Multivariate Analysis of Variance (PERMANOVA) using Distance Matrices. Although not consistently correlating with univariate measures, the ACI was significantly interacting with the changing benthic assemblage composition, as it changed over time and protection (Inside vs Outside the MPA).

ACI showed potential to allude to shifting benthic communities, yet with no consistency when used alongside univariate measures of diversity. Although it is not without its own disadvantages, and thus should be developed further before implementation, the ACI could potentially reflect more complex changes to the benthos than simply the overall diversity.

Keywords: Acoustic Complexity Index, Biodiversity, Marine Protected Area, Monitoring Tools

1. Introduction

Biodiversity provides a useful measure to assess ecosystem health (Worm et al., 2006), and is increasingly being used for conservation and monitoring purposes, with an observed decrease used as a proxy for a degraded or negatively impacted ecosystem (Wabnitz et al., 2018). To quantify and

compare these changes in diversity, many univariate indices have been produced, which simplify an assemblage of taxa into a single value. The most commonly used indices involve integrating the number of species present with measures of how the species are distributed within the assemblages, such as Number of Species (Kaplan et al., 2015; Pieretti and Farina, 2013; Sheehan et al., 2013b), Shannon-Wiener's diversity index (De-La-Ossa-Carretero et al., 2012), Simpson's diversity index (Miralles et al., 2016; Rombouts et al., 2019) and taxonomic distinctness (Clarke and Warwick, 2001; Leonard et al., 2006), which also involves phylogenetic distance.

Historic methods for assessing marine biodiversity have often used destructive practices (Francour, 1994; Lipej et al., 2003), such as poisoning (Diamant et al., 1986) or trawling (Cappo et al., 2004). However, for the study of recovering and fragile benthic systems, such as those in Marine Protected Areas (MPAs), non-invasive, non-extractive methods such as Underwater Visual Census (UVC) or Underwater Video Survey (UVS) are considered more appropriate (Sheehan et al., 2013a, 2010). Visual methods will always have the drawback that there is no physical sample taken, although image libraries give a permanent record, and thus those species that are harder to identify visually will always be under-sampled; yet this lack of physical sample means the populations being researched are almost or completely unaffected by the survey taken. A potential addition to supplement visual survey would be the assessment of the marine soundscape (Staaterman et al., 2017). This method for sampling the marine environment is similarly non-extractive and non-invasive, while sampling components of the ecosystem potentially under-represented by visual methods alone.

The marine soundscape comprises both natural and anthropogenic elements. Assessment of the biological element (biophony) of the marine soundscape has been used to describe overall biodiversity (Bertucci et al., 2016), reproductive behaviour (de Jong et al., 2018), habitat selection (Vermeij et al., 2010), spawning (Casaretto et al., 2014; Hawkins and Amorim, 2000) and predator-prey interactions (Bernasconi et al., 2011; Giorli et al., 2016). Biophony is produced by a wide range of taxa ranging from large cetaceans producing low frequency (~20Hz) calls or songs (Samaran et al., 2013), that can be detected up to thousands of kilometres away (Rivers, 1997), to crustaceans creating loud (190 dB re 1 μ Pa), broadband (2kHz up to 300kHz) 'snaps' and 'pops' (Picciulin et al., 2013).

Acoustic indices have been developed and utilised in marine (Gordon et al., 2018; Harris et al., 2016; Nedelec et al., 2015; Pieretti et al., 2017; Trenkel et al., 2011) and terrestrial (Farina and Pieretti, 2014; Merchant et al., 2015; Pieretti et al., 2015, 2011; Pijanowski et al., 2011; Sueur et al., 2008b) environments to assess whole ecosystem biodiversity. The use of these acoustic indices is perceived to allow hidden or shy species, overlooked by other survey methods, to be accounted for (Staaterman et al., 2017). The ACI as set out in Pieretti *et al.* (2011) quantifies the relative change in sound intensity across all frequencies of a soundscape, while being minimally affected by constant anthropogenic noise. The ACI was developed on the assumption that with increased diversity of species, there would be an increase in the complexity of biological sound produced. So far, most analyses of ACI have shown a positive correlation with a variety of biodiversity indices (Bertucci et al., 2016; Harris et al., 2016; Meyer et al., 2018; Pieretti et al., 2015, 2011).

The two survey methods, visual and acoustic, are thought to complement each other by overlapping, as well as covering differing spatial scales and taxonomic groups (Staaterman et al., 2017). However, the majority of studies to date regarding this interaction have been based either in areas of very high biodiversity, such as coral reef systems (Bertucci et al., 2016; Kaplan et al., 2015), or only focused on fish diversity (Harris et al., 2016). As such, the transferability to other habitats and ecosystems is limited.

This study assessed the suitability of the ACI index derived from using acoustic recording as a monitoring method and to explore its relationship with seabed biodiversity. As such, a 5 year study within a recovering temperate reef seabed ecosystem was undertaken, in which were protected areas and those open to bottom fishing.

It was expected that the ACI and two visual biodiversity indices, Number of Species and Shannon's Diversity Index, derived from Baited Remote Underwater Video systems (BRUVs) data ('visual biodiversity indices' from now on), would increase over time in the MPA relative to the areas that continue to be fished. As a recovering system it would be predicted that the interaction of time and treatment would be significant. Therefore, the following hypotheses were assessed for inside vs outside the MPA:

1. The ACI would increase over time,
2. The visual biodiversity indices would increase over time,
3. The visual biodiversity indices and the ACI would correlate with each other over time,
4. Changes in the mobile benthic assemblage composition would result in similar changes to the ACI.

2. Methods

2.1 Study Location

Lyme Bay (Fig. 1), is located on the south coast of England, and contains areas of rocky reef habitat known to include nationally important fragile reef building species (Hiscock and Breckels, 2007). A Statutory Instrument (SI), a type of MPA, was established in 2008 in Lyme Bay. The SI excluded all towed demersal fishing equipment (scallop dredging and trawling) from a 206 km² area of the bay.

Experimental site selection was based on similar biotope classifications to negate any confounding effects of habitat heterogeneity (Claudet et al., 2008), with all sites being on either hard or ‘mixed’ substrate at similar depths (Sheehan et al., 2013b; Stevens et al., 2014). There were two treatments: Inside the MPA (n=12) and Outside the MPA (n=6). Geographically similar pairs of sites were grouped into ‘Areas’.

In the winter of 2013/2014, the south coast of the UK experienced severe and unprecedented storm activity that was observed to have major impacts on South West England's coastal systems (Masselink et al., 2016). The effect to both the protected and non-protected ecosystems provided an opportunity to start a new monitoring strategy. This incorporated acoustic recording and assessment of the marine soundscape inside and outside of the MPA alongside visual measures of the seabed assemblage and allowed the assessment of the emerging acoustic analyses.




Fig. 1. Map of the UK inlaid in a map of Lyme Bay, showing site locations and their treatments (Blue Cross: Marine Protected Area, Red Circle: Open Control). Dark blue line shows the Statutory Instrument and light blue lines show 10m depth contours.

2.2 Data Collection

2.2.1 Acoustic Recorder Deployment

At each site, an acoustic recorder was attached and deployed with one of the three replicate BRUVs (Fig. 2). The acoustic recorders used were low power Digital SpectroGrams (DSG) (Hydrophone Calibration Sensitivity=-190dBV/uPa, Sample rate=50 kHz, Decimation Factor=4, System Gain=20, Effective Sample Range=0-25 kHz; Loggerhead Instruments, Sarasota, FL, USA), which were used to record DSG files on a duty cycle of 16 seconds recorded every 2 minutes to conserve battery life. The recorders were attached to one BRUVs for every site, to sample identical locations (Fig. 2), but, as

DSG acoustic recorders sample a larger area (Simard *et al.*, 2015) than the BRUVs maximum effective range of attraction (AR) (Cappo *et al.*, 2004), single acoustic recordings were used across the three BRUVs replicates (Fig. S1).




Fig. 2. Sampling design for BRUVs and Acoustic Recorders.

2.2.2 Acoustic File Extraction

For each deployment, audio recordings were prepared to ~360 second WAV format files, equating to 45 minutes, to align with the recorded BRUVs. Auditory and visual examinations were then used to remove any sporadic dominant anthropogenic interference using ‘seewave’ package in R (Sueur *et al.*, 2008a). For each deployment Acoustic Complexity Index (ACI)(Pieretti *et al.*, 2011) was calculated, using the R packages ‘tuneR’ and ‘seewave’ (Ligges *et al.*, 2016; Sueur *et al.*, 2008a).

2.2.3 Acoustic Complexity Index

Originally developed to analyse terrestrial avian communities, the Acoustic Complexity Index (ACI) quantifies the change in adjacent spectrogram intensities for all temporal steps and frequency bins of a recording (Pieretti *et al.*, 2011). Firstly, sound files were split into frequency bins and temporal steps. The change in adjacent intensities are then summed across these frequency bins and temporal steps. Thus, high ACI values are produced by large variations in sound intensity across many frequencies and times, whereas constant levels of similar intensity, such as most anthropogenic sources (e.g. boat engine), will produce low values of ACI (Bertucci *et al.*, 2016).

ACI was chosen for the current study since the hydrophones used were encased in resin and could not be calibrated, which is a necessary requirement for calculating amplitude. As such, the acoustic files created could not be analysed with other popular acoustic measures which rely on amplitude, such as

Acoustic Entropy, Acoustic Richness, Root Mean Square or Sound Pressure Level (Picciulin et al., 2013; Sueur et al., 2008b).

2.3 Baited Remote Underwater Video Systems

Baited Remote Underwater Video systems (BRUVs) are a non-destructive method for sampling mobile communities (Babcock et al., 1999; Heagney et al., 2007). Three replicate BRUVs were deployed at each site (Fig. 2) for 45 minutes then recovered. Specifications of equipment are described in Bicknell *et al.*, (2019).

2.3.1 Video analysis

After a preliminary settling period of 5 minutes, 30 minutes of video were analysed in 1 minute segments. For each segment all mobile benthic organisms were identified and recorded. All organisms were identified to the highest taxonomic resolution possible. Abundance (MaxN) was calculated for each species from the maximum number of individuals of each species observed across all of the 30 minute segments.

2.4 Statistical Analysis

Permutational Analysis of Variance (PERMANOVA) was used to test differences in between years and treatments for the ACI, Shannon's Diversity Index, Number of Species and the assemblage composition. Year and Treatment were fixed factors with five and two levels respectively (Year: 2014, 2015, 2016, 2017 and 2018; Treatment: MPA and Open Control) using Primer v7 and PERMANOVA+ (Anderson et al., 2008; Clarke and Gorley, 2015). The assemblage composition analysis also included a random factor Area, which was nested inside Treatment (MPA=6 areas, OC=3 areas). PERMANOVA was chosen as it is robust to unbalanced designs (Sheehan et al., 2013b). For Shannon's Index, Number of Species and assemblage composition, the ACI was included as a covariate. The statistical significance of the variance components were tested using 9999 permutations under a reduced model (Anderson, 2001; Anderson and ter Braak, 2002). The analyses of the ACI and the two visual biodiversity indices were undertaken on the basis of a Euclidean distance matrix calculated from the Index values (Anderson and Millar, 2004). The assemblage

composition analysis was based on a Bray-Curtis dissimilarity matrix calculated from dispersion weighted, fourth root transformed abundance data. Significant interactions ($p < 0.05$) of fixed terms were tested using PERMANOVA pairwise tests.

To assess correlations between visual biodiversity measures and the ACI, scatter plots were created with Pearson correlations showing R values and significance ($p < 0.05$). Assemblage composition was visualised using non-metric Multi-Dimensional Scaling (nMDS: Clarke and Gorley, 2015).

3. Results

3.1 Acoustic Complexity Index

The interaction between year and treatment was significant for the ACI (Table 1: Pseudo-F=2.6766, $p = 0.0351$). This significant interaction shows that there is a combined effect of year and treatment.

The MPA was more acoustically complex than Open Controls (OC) in 2014 and 2018 (Table 1; 2014: $p = 0.009$; 2018: $p = 0.0288$), whereas the OC group was more complex in 2016 (Fig. 3A, Table 1; 2016: $p = 0.0218$). Overall across all years, mean ACI was lower inside the MPA (1.4% lower than outside: Fig. 3A).

*Table 1 Results table of PERMANOVA analysis of Euclidean distances assessing Acoustic Complexity Index with Year and Treatment as interactions and Pairwise comparisons of yearly differences between Treatments for the ACI. (Bold p values denotes significance, *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$).*

Source	PERMANOVA				Pairwise Comparisons		
	df	MS	Pseudo-F	p	Year	MPA vs OC	
						t	p
Year	4	849.5	8.6736	0.0001***	2014	2.7956	0.009**
Treatment	1	255.76	2.6113	0.1117	2015	0.33271	0.9741
Year x Treatment	4	262.15	2.6766	0.0351*	2016	2.3627	0.0218*
					2017	1.6034	0.1182

Residual	168	97.941	2018	2.2848	0.0288*
-----------------	-----	--------	-------------	--------	----------------

3.2 Visual Biodiversity Indices

Both indices of diversity (Shannon's Diversity Index and Number of Species) were greater on average inside vs outside the MPA, with a mean percentage difference from outside to inside of: 20.0% for Shannon's and 8.4% for Number of Species (Fig. 3B & 3C).

When all diversity indices, both acoustic and visual biodiversity, are analysed by year within treatment, there is no significant trend with year displayed by the ACI, either inside or outside the MPA (Fig. 4, Inside: $R=0.14$, $p>0.05$; Outside $R=0.18$, $p>0.05$). However, Number of Species significantly increased with time both inside and outside the MPA (Fig. 4: Inside: $R=0.36$, $p<0.0001$, Outside: $R=0.32$, $p=0.02$). Outside the MPA, Shannon's index shows no significant trend with time and has a small significant increase with time inside the MPA (Fig. 4: Inside: $R=0.21$, $p=0.017$; Outside: $R=5.7 \times 10^{-4}$, $p>0.05$).




Fig. 3. Mean \pm SE Acoustic Complexity (A), Number of Species (B) and Shannon's Diversity (C) Inside (Filled Blue) and Outside the MPA (Unfilled Red) across all years.




Fig. 4. Scatter plot with Pearson correlation coefficient for Year against Acoustic Complexity Index (top), Number of Species (middle) and Shannon's Diversity Index (bottom) split by treatment (Inside: blue and left, Outside: red and right). Includes R values and significance shown by ns: $p > 0.05$, *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$. Shading around regression line shows 95% confidence interval.

3.3 Relationship between Visual Biodiversity and Acoustic Complexity

Shannon's Index was greater in the MPA than the Open Controls but the relationship with Year was marginally non-significant (Table 2; Treatment: Pseudo- $F=10.726$, $p=0.0013$; Year: Pseudo- $F=2.3123$, $p=0.0564$). For the Number of Species there was a significant interaction between the Year and the ACI (Pseudo- $F=6.4837$, $p=0.0002$) as well as ACI and Treatment (Pseudo- $F=6.1875$, $p=0.0157$). This shows that, although not correlating with Number of Species, the ACI is interacting when the factors Year and Treatment are introduced. However, this is not significant for the Year x Treatment interaction.

Table 2. Results table of PERMANOVA analysis on Euclidean distance assessing Shannon's Diversity Index (A) and Number of Species (B) with Year and Treatment as interactions and ACI as a covariate. (Bold p values denotes significance, *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$).

Source	df	Shannon's Diversity Index			Number of Species		
		MS	Pseudo-F	p	MS	Pseudo-F	p
ACI	1	0.17249	0.89629	0.3526	1.5537	0.34091	0.5553
Year	4	0.445	2.3123	0.0564	53.05	11.64	0.0001***
Treatment	1	2.0641	10.726	0.0013 **	6.3986	1.4039	0.2415
ACI x Year	4	0.30352	1.5771	0.1812	29.55	6.4837	0.0002**
ACI x Treatment	1	0.21135	1.0982	0.2936	28.2	6.1875	0.0157*
Year x Treatment	4	0.25056	1.3019	0.2712	10.249	2.2487	0.0635
ACI x Year x Treatment	4	0.27228	1.4148	0.2248	5.133	1.1262	0.3408
Residuals	158	0.19245			4.5576		

Neither Number of Species nor Shannon's Index correlated with the ACI when compared across all the years and treatments (Fig. 5; Shannon's Index: $R = -0.37$, $p > 0.05$ and Number of Species: $R = -0.067$, $p > 0.05$). However, within each year, Number of Species did correlate with ACI with the exception of 2016. However, the orientation of this correlation was inconsistent; it was positive for 2014 and 2018, and negative for 2015 and 2017 (Fig. 6; Positive- 2014: $R = 0.36$, $p = 0.041$; 2018: $R = 0.4$, $p = 0.041$; Negative- 2015: $R = -0.42$, $p = 0.017$; 2017: $R = -0.57$, $p < 0.001$). In contrast, Shannon's Index correlated with ACI in only 2017; this correlation was negative (Fig. 6; 2017: $R = -0.4$, $p = 0.017$).




Fig. 5 Scatter plot with Pearson correlation coefficient for Acoustic Complexity against Number of Species (above) and Shannon's Diversity Index (below). R values are shown and significance shown by ns: $p > 0.05$, *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$. Shading around regression line shows 95% confidence interval.




Fig. 6. Scatter plot with Pearson correlation coefficient for Number of Species (left) and Shannon's Diversity Index (right) against Acoustic Complexity Index. R values are shown and significance shown by ns: $p > 0.05$, *: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$. Shading around regression line shows 95% confidence interval.

3.4 Mobile Benthic Assemblage Composition

The assemblage compositions of the two treatments diverged with increasing time with the two treatments changing at different rates (Fig. 7; Table 3: ACI x Year x Treatment: Pseudo-F: 1.7682, $p=0.0482$). Pearson correlation of more than 0.85 showed the reptant decapod crustaceans *Inachus* spp. and *Pagurus* spp. were most important to the Open Control composition (Fig.7), whereas, the species most important for the MPA assemblage were the wrasse species *Labrus mixtus*, *Labrus bergylta* and *Ctenolabrus rupestris* (Fig. 7).




Fig. 7. Non-Metric Multidimensional Scaling plot of distance to centroids split by Year and Treatment from adjusted Bray-Curtis similarity of fourth root transformed abundance data. Points are labelled by Year and coloured by treatment (blue: Inside MPA, red: Outside MPA) and scaled according to mean ACI values. Vectors overlaid display 0.85 Pearson correlation for the species driving the difference in the assemblage composition.

Table 3. Results table of PERMANOVA analysis on adjusted Bray-Curtis similarity assessing mobile benthic assemblage composition with Year and Treatment as fixed interactions, area as a random interaction nested within treatment and ACI as a covariate. (Bold p values denotes significance, *: $p<0.05$, **: $p<0.01$, ***: $p<0.001$).

Source	Assemblage Composition			
	df	MS	Pseudo-F	p
ACI	1	4003.3	2.0845	0.0457*
Year	4	9011.3	3.4253	0.0001***

Treatment	1	29580	3.652	0.0003***
Area(Treatment)	7	8302.8	8.974	0.0001***
ACI x Year	4	2167.7	1.5326	0.0607
ACI x Treatment	1	2296.2	2.4819	0.0177*
Year x Treatment	4	1883.9	1.1026	0.3666
ACI x Area(Treatment)	7	1338.7	1.4469	0.0328*
Year x Area(Treatment)	23	1419.6	1.5344	0.0001***
ACI x Year x Treatment	2	1635.9	1.7682	0.0482*
Residuals	123	925.2		

4. Discussion

After high storm activity impacted the coastal systems of Lyme Bay and beyond (Masselink et al., 2016), acoustic and BRUV monitoring was implemented. It was hypothesised that the Acoustic Complexity Index would increase over time as the biodiversity of the area increased. Furthermore, the ACI was expected to be greater inside the protected area in comparison to the surrounding fished areas. Finally it was hypothesised that the ACI would change in a similar pattern to that of the mobile benthic assemblage composition recorded by BRUV systems.

4.1 Visual Biodiversity and Acoustic Indices

The Acoustic Complexity Index, as a covariate, showed a greater number of significant interactions alongside treatment and year with Number of Species than Shannon's Diversity (Fig. 4 & 6; Table 2). This relationship between ACI and Number of Species implies that the ACI is less affected by abundance and more by the number of species present. This is to be expected as the ACI was developed under the theory that many differing biological noises in an environment imply many different species (Pieretti et al., 2011). Yet, when studying fish vocal communities specifically, the ACI shows little discrimination between the abundance and the diversity of sound (Bolgan et al., 2018). Thus, the features that the ACI is enumerating from the marine soundscape may not, as hypothesised, correspond directly to the species detected by the BRUVs represented in the biodiversity indices tested. This may explain the yearly correlations between ACI and Number of Species, which changed orientation (between positive and negative), while also displaying small

effect sizes throughout the study. This complete reversal, at times, could be misleading if studies using this method do not cover an appropriate temporal scale. This inconsistency may be the result of abiotic or anthropogenic noises (McWilliam and Hawkins, 2013), or specific species and behaviours dominating the soundscape, meaning the presence of specific species in the acoustically sampled area, but not recorded by the BRUVs, could be driving this inconsistent pattern and therefore may preclude it from certain applications, such as directly replacing more traditional biodiversity monitoring methods.

4.2 Shifting Benthic Composition

There was a clear divergence of assemblage compositions, between inside and outside treatment areas moving further apart year on year (Fig. 7). However, without data on the ‘before fishing’ assemblage, it would be very difficult to suggest whether this separation is recovery of the ecosystem. Yet, Pearson’s correlations would suggest that the species most associated with the MPA are classed as reef dwelling species: *Ctenolabrus rupestris* remain in the same local area for several years, thus, maintaining their ‘territory’ (Darwall et al., 1992). The Open Controls were dominated by the scavenging species *Inachus* spp. and *Pagurus* spp. (Fig. 7): both have broad habitat preferences although *Inachus* spp. is more likely to be found on mixed coarse substrata (Rowley, 2008).

Although not significant, inside the MPA there was a higher Number of Species and Shannon’s Diversity (Fig. 3). Both indices increasing with treatment (Shannon’s Index) and year (Number of Species) show that the MPA in Lyme Bay acts as a refuge to allow biodiversity to increase (Fig. 4: Table 2). The assemblage composition does not interact with changing ACI alone, but is significant when aligned with year and treatment, which would be expected of a recovering system (Table 3). This would suggest that, although it did not correlate overall with visual biodiversity, the ACI is sensitive to some level of the non-background variation in the assemblage composition.

4.3 Applications of the Acoustic Complexity Index

Research into acoustic recordings is such a growing area due to its ability to provide information on local assemblage structure (Pijanowski et al., 2011; Sueur and Farina, 2015). Yet, as with most areas

of ecology, the transition from the terrestrial to the marine poses a new range of obstacles to overcome (Giorli, 2016; Radford et al., 2011; Ricci et al., 2017). Many different indices have been produced to quantify marine biological processes, such as Acoustic Richness, Acoustic Entropy Index and Acoustic Complexity Index (Gage and Axel, 2014; Lillis et al., 2014; McWilliam and Hawkins, 2013; Staaterman et al., 2014). Their use as proxies for marine biodiversity has been assessed (Harris et al., 2016), with the Acoustic Complexity Index being the most favoured (Lindseth and Lobel, 2018) both alone and in combination with other acoustic indices (Gordon et al., 2018).

The Acoustic Complexity Index has been shown to have a number of drawbacks (Kaplan et al., 2015; McWilliam and Hawkins, 2013). These drawbacks can arise from interference by the biophony, geophony or anthrophony. For example, the ACI has shown to be increased heavily by snapping shrimp, which produce a high intensity broadband ‘snap’, meaning an increased ACI when diversity has only marginally increased (McWilliam and Hawkins, 2013). In contrast, chorusing behaviour can heavily decrease ACI (Kaplan et al., 2015). Hence, ACI in certain situations can be dominated by either few or many species, producing opposing changes in the ACI and the observed biodiversity. The assemblage composition outside of the MPA, in this case, was heavily dominated by hermit crabs of the genus *Pagurus*. It is possible that these large aggregations of *Pagurus* spp. (up to 70 in one video), which ‘rap’ on others’ shells for shell competition (Edmonds and Briffa, 2016), dominated the ACI in a similar way to snapping shrimps. Dominance of snapping shrimp in the marine soundscape affects most other acoustic indices, not just the ACI (Au and Banks, 1998; Lindseth and Lobel, 2018; Radford et al., 2008). Thus, this issue needs to be overcome for multiple different methods. The ACI can also be heavily influenced by geophony such as wind and rain (McWilliam and Hawkins, 2013). Although designed to minimise the influence of anthropogenic inputs, ACI will also be affected by any sounds which are not repetitive or consistent in intensity sounds, such as boat engines (Pieretti et al., 2011). Therefore, post sampling examination of the recordings was carried out here to minimise any sporadic dominant abiotic or anthropogenic interference, which would otherwise influence the ACI.

All recordings here were made during the day and, as such, potentially not at the highest acoustic activity times, which for most fish are dawn and dusk (Bertucci et al., 2017, 2016, 2015; Radford et al., 2014). Further investigation into this index should include diurnal recording strategies, while also taking into consideration the activity cycles based upon lunar phase (Harris et al., 2016; Staaterman et al., 2014). As shown here, correlations between the ACI and other diversity measures can occur, but can vary considerably in their orientations over years. Thus, temporal scales which include lunar and daily cycles, should be used to assess these indices. Although not possible here, the combination of multiple metrics together has been suggested to provide a more robust assessment of the marine soundscape (Gordon et al., 2018). However, the individual aspects of the soundscape which ACI quantifies need to be further understood before it can be appropriately combined with other metrics.

The use of ACI in this MPA, off the south coast of the UK, has not shown the direct relationship with the observed ecology as demonstrated elsewhere (Harris et al., 2016; Picciulin et al., 2013), yet did show significant interactions across treatments and years. As the significant interactions were found under multivariate and not univariate analysis, it is likely that the ACI is quantifying other elements of the marine soundscape and not just the diversity of species (Bolgen et al., 2018; Kaplan et al., 2015; McWilliam and Hawkins, 2013). For this, or another, acoustic index to be used as a rapid and cost-effective monitoring tool, the drawbacks mentioned here need to be addressed. Yet more importantly, the elements of the marine soundscape, which the ACI is quantifying, need to be better understood. Subsequently, thorough experimental assessments will be needed, with robust spatial and temporal coverage. This is essential, as based on a single year of this study (e.g. 2014 or 2018), ACI would have shown a positive correlation with Number of Species that has been found elsewhere. Again, this shows that temporal and geographical scales are important considerations for the development of any such index or method.

4.4 Conclusions

In conclusion, the Acoustic Complexity Index is not as yet ready to be used as a standalone marine diversity monitoring tool. In conjunction with other methods, such as BRUVs, which showed the recovery and increased diversity within the Lyme Bay MPA, this acoustic index shows potential to

allude to shifting benthic assemblage compositions. Yet this was not seen with consistency when used alongside univariate measures of diversity. This implies that although it is not without its own disadvantages, the ACI is demonstrating more complex changes than overall univariate diversity. This potential as a tool for rapidly assessing a large area of the marine environment makes it highly attractive. However, for it to be used as a monitoring tool, the information it provides regarding shifting assemblage compositions and diversity needs to be fully researched and understood.

Acknowledgements

To carry out fieldwork thanks are given to Lyme Bay Fishers John Walker, Robert King and Keiran Perree and University of Plymouth staff and student volunteers especially Amy Cartwright for her fieldwork support and logistics. Also, thank you to Marti Anderson for advice regarding statistical analysis. Funding: This work was supported by Natural England and The European Commission [EMFF RETURN ENG1388].

Authors' Contributions

EVS and MJA conceived the ideas and monitoring design; MJW provided technical advice regarding acoustic analytical methods; EVS, LH, AR and BFRD collected data; BFRD and LH organized and analysed data; BFRD, EVS and LH led the writing of the manuscript. All authors contributed critically to drafts and gave final approval for publication.

References:

- Anderson, M.J., 2001. A new method for non-parametric multivariate analysis of variance. *Austral Ecol.* 26, 32–46. <https://doi.org/10.1046/j.1442-9993.2001.01070.x>
- Anderson, M.J., Gorley, R.N., Clarke, K.R., 2008. PERMANOVA+ for PRIMER: Guide to Software and Statistical Methods.
- Anderson, M.J., Millar, R.B., 2004. Spatial variation and effects of habitat on temperate reef fish assemblages in northeastern New Zealand. *J. Exp. Mar. Bio. Ecol.*

355 <https://doi.org/10.1016/j.jembe.2003.12.011>

356 Anderson, M.J., ter Braak, C.J.F., 2002. Permutation tests for multi-factorial analysis of variance. J.
 357 Stat. Comput. Simul. 73, 85–113.

358 Au, W.W.L., Banks, K., 1998. The acoustics of the snapping shrimp *Synalpheus parneomeris* in
 359 Kaneohe Bay. J. Acoust. Soc. Am. 103, 41. <https://doi.org/10.1121/1.423234>

360 Babcock, R.C., Kelly, S., Shears, N.T., Walker, J.W., Willis, T.J., 1999. Changes in community structure
 361 in temperate marine reserves. Mar. Ecol. Prog. Ser. 189, 125–134.

362 Bernasconi, M., Nøttestad, L., Axelsen, B.E., Krakstad, J.-O., 2011. Acoustic observations of dusky
 363 dolphins *Lagenorhynchus obscurus* hunting Cape horse mackerel *Trachurus capensis* off
 364 Namibia. Mar. Ecol. Prog. Ser. 429, 209–218. <https://doi.org/10.3354/meps09071>

365 Bertucci, F., Parmentier, E., Berten, L., Brooker, R.M., Lecchini, D., 2015. Temporal and Spatial
 366 Comparisons of Underwater Sound Signatures of Different Reef Habitats in Moorea Island,
 367 French Polynesia. PLoS One 10, e0135733. <https://doi.org/10.1371/journal.pone.0135733>

368 Bertucci, F., Parmentier, E., Berthe, C., Besson, M., Hawkins, A.D., Aubin, T., Lecchini, D., 2017.
 369 Snapshot recordings provide a first description of the acoustic signatures of deeper habitats
 370 adjacent to coral reefs of Moorea. PeerJ 5, e4019. <https://doi.org/10.7717/peerj.4019>

371 Bertucci, F., Parmentier, E., Lecellier, G., Hawkins, A.D., Lecchini, D., 2016. Acoustic indices provide
 372 information on the status of coral reefs: An example from Moorea Island in the South Pacific.
 373 Sci. Rep. 6, 1–9. <https://doi.org/10.1038/srep33326>

374 Bicknell, A.W.J., Sheehan, E. V., Godley, B.J., Doherty, P.D., Witt, M.J., 2019. Assessing the impact of
 375 introduced infrastructure at sea with cameras: A case study for spatial scale, time and
 376 statistical power. Mar. Environ. Res. 147, 126–137.
 377 <https://doi.org/10.1016/j.marenvres.2019.04.007>

378 Bolgan, M., Amorim, M.C.P., Fonseca, P.J., Iorio, L. Di, Parmentier, E., 2018. Acoustic Complexity of
 379 vocal fish communities: a field and controlled validation. *Sci. Rep.* 8, 10559.
 380 <https://doi.org/10.1038/s41598-018-28771-6>

381 Cappo, M., Speare, P., De'ath, G., 2004. Comparison of baited remote underwater video stations
 382 (BRUVS) and prawn (shrimp) trawls for assessments of fish biodiversity in inter-reefal areas of
 383 the Great Barrier Reef Marine Park. *J. Exp. Mar. Bio. Ecol.* 302, 123–152.
 384 <https://doi.org/10.1016/j.jembe.2003.10.006>

385 Casaretto, L., Picciulin, M., Olsen, K., Hawkins, A.D., 2014. Locating spawning haddock
 386 (*Melanogrammus aeglefinus*, Linnaeus, 1758) at sea by means of sound. *Fish. Res.* 154, 127–
 387 134. <https://doi.org/10.1016/j.fishres.2014.02.010>

388 Clarke, K.R., Gorley, R.N., 2015. *PRIMER v7: User Manual/Tutorial*.

389 Clarke, K.R., Warwick, R.M., 2001. A further biodiversity index applicable to species lists: Variation in
 390 taxonomic distinctness. *Mar. Ecol. Prog. Ser.* 216, 265–278.
 391 <https://doi.org/10.3354/meps216265>

392 Claudet, J., Osenberg, C.W., Benedetti-Cecchi, L., Domenici, P., García-Charton, J.A., Pérez-Ruzafa, Á.,
 393 Badalamenti, F., Bayle-Sempere, J.T., Brito, A., Bulleri, F., Culioli, J.M., Dimech, M., Falcón, J.M.,
 394 Guala, I., Milazzo, M., Sánchez-Meca, J., Somerfield, P.J., Stobart, B., Vandeperre, F., Valle, C.,
 395 Planes, S., 2008. Marine reserves: Size and age do matter. *Ecol. Lett.* 11, 481–489.
 396 <https://doi.org/10.1111/j.1461-0248.2008.01166.x>

397 Darwall, W.R.T., Costello, M.J., Donnelly, R., Lysaght, S., 1992. Implications of life-history strategies
 398 for a new wrasse fishery. *J. Fish Biol.* 41, 111–123. [https://doi.org/10.1111/j.1095-](https://doi.org/10.1111/j.1095-8649.1992.tb03873.x)
 399 [8649.1992.tb03873.x](https://doi.org/10.1111/j.1095-8649.1992.tb03873.x)

400 De-La-Ossa-Carretero, J.A., Del-Pilar-Ruso, Y., Giménez-Casaldueiro, F., Sánchez-Lizaso, J.L., Dauvin,
 401 J.C., 2012. Sensitivity of amphipods to sewage pollution. *Estuar. Coast. Shelf Sci.* 96, 129–138.

402 <https://doi.org/10.1016/j.ecss.2011.10.020>

403 de Jong, K., Amorim, M.C.P., Fonseca, P.J., Fox, C.J., Heubel, K.U., 2018. Noise can affect acoustic
404 communication and subsequent spawning success in fish. *Environ. Pollut.* 237, 814–823.
405 <https://doi.org/10.1016/j.envpol.2017.11.003>

406 Diamant, A., Tuvia, A. Ben, Baranes, A., Golani, D., 1986. An analysis of rocky coastal eastern
407 Mediterranean fish assemblages and a comparison with an adjacent small artificial reef. *J. Exp.*
408 *Mar. Bio. Ecol.* 97, 269–285. [https://doi.org/10.1016/0022-0981\(86\)90245-5](https://doi.org/10.1016/0022-0981(86)90245-5)

409 Edmonds, E., Briffa, M., 2016. Animal behaviour Weak rappers rock more: hermit crabs assess their
410 own agonistic behaviour. *Biol. Lett.* 12, 20150884. <https://doi.org/10.1098/rsbl.2015.0884>

411 Farina, A., Pieretti, N., 2014. Sonic environment and vegetation structure: A methodological
412 approach for a soundscape analysis of a Mediterranean maqui. *Ecol. Inform.*
413 <https://doi.org/10.1016/j.ecoinf.2013.10.008>

414 Francour, P., 1994. Pluriannual analysis of the reserve effect on ichthyofauna in the Scandola natural
415 reserve (Corsica , Northwestem Mediterranean). *Oceanol. Acta* 17, 309–317.

416 Gage, S.H., Axel, A.C., 2014. Visualization of temporal change in soundscape power of a Michigan
417 lake habitat over a 4-year period. *Ecol. Inform.* 21, 100–109.
418 <https://doi.org/10.1016/j.ecoinf.2013.11.004>

419 Giorli, G., 2016. Temporal and spatial variation of beaked and sperm whales foraging activity in
420 Hawai’i, as determined with passive acoustics. *J. Acoust. Soc. Am.* 140, 2333.
421 <https://doi.org/10.1121/1.4964105>

422 Giorli, G., Neuheimer, A., Copeland, A., Au, W.W.L., 2016. Temporal and spatial variation of beaked
423 and sperm whales foraging activity in Hawai’i, as determined with passive acoustics. *J. Acoust.*
424 *Soc. Am.* 140, 2333–2343. <https://doi.org/10.1121/1.4964105>

425 Gordon, T.A.C., Harding, H.R., Wong, K.E., Merchant, N.D., Meekan, M.G., McCormick, M.I., Radford,
 426 A.N., Simpson, S.D., 2018. Habitat degradation negatively affects auditory settlement behavior
 427 of coral reef fishes. *Proc. Natl. Acad. Sci.*

428 Harris, S.A., Shears, N.T., Radford, C.A., 2016. Ecoacoustic indices as proxies for biodiversity on
 429 temperate reefs. *Methods Ecol. Evol.* 7, 713–724. <https://doi.org/10.1111/2041-210X.12527>

430 Hawkins, A.D., Amorim, M.C.P., 2000. Spawning Sounds of the Male Haddock, *Melanogrammus*
 431 *aeglefinus*. *Environ. Biol. Fishes* 59, 29–41.

432 Heagney, E.C., Lynch, T.P., Babcock, R.C., Suthers, I.M., 2007. Pelagic fish assemblages assessed using
 433 mid-water baited video: standardising fish counts using bait plume size. *Mar. Ecol. Prog. Ser.*
 434 350, 255–266. <https://doi.org/10.3354/meps07193>

435 Hiscock, K., Breckels, M., 2007. Marine Biodiversity Hotspots in the UK A report identifying and
 436 protecting areas for marine biodiversity Marine Biodiversity Hotspots in the UK: their
 437 identification and protection. WWF.

438 Kaplan, M.B., Mooney, T.A., Partan, J., Solow, A.R., 2015. Coral reef species assemblages are
 439 associated with ambient soundscapes. *Mar. Ecol. Prog. Ser.* 533, 93–107.
 440 <https://doi.org/10.3354/meps11382>

441 Leonard, D.R.P., Clarke, K.R., Somerfield, P.J., Warwick, R.M., 2006. The application of an indicator
 442 based on taxonomic distinctness for UK marine biodiversity assessments. *J. Environ. Manage.*
 443 78, 52–62. <https://doi.org/10.1016/j.jenvman.2005.04.008>

444 Ligges, U., Krey, S., Mersmann, O., Schnackenberg, S., 2016. tuneR: Analysis of music. URL:[http://r-](http://r-forge.r-project.org/projects/tuner/)
 445 [forge.r-project.org/projects/tuner/](http://r-forge.r-project.org/projects/tuner/).

446 Lillis, A., Eggleston, D.B., Bohnenstiehl, D.R., 2014. Estuarine soundscapes: distinct acoustic
 447 characteristics of oyster reefs compared to soft-bottom habitats. *Mar. Ecol. Prog. Ser.* 505, 1–
 448 17. <https://doi.org/10.3354/meps10805>

449 Lindseth, A., Lobel, P., 2018. Underwater Soundscape Monitoring and Fish Bioacoustics: A Review.
 450 Fishes 3, 36. <https://doi.org/10.3390/fishes3030036>

451 Lipej, L., Bonaca, M.O., Sea, A., 2003. Coastal Fish Diversity in Three Marine Protected Areas and One
 452 Unprotected Area in the Gulf of Trieste (Northern Adriatic). *Mar. Ecol.* 24, 259–273.

453 Masselink, G., Scott, T., Poate, T., Russell, P., Davidson, M., Conley, D., 2016. The extreme 2013/2014
 454 winter storms: Hydrodynamic forcing and coastal response along the southwest coast of
 455 England. *Earth Surf. Process. Landforms* 41, 378–391. <https://doi.org/10.1002/esp.3836>

456 McWilliam, J.N., Hawkins, A.D., 2013. A comparison of inshore marine soundscapes. *J. Exp. Mar. Bio.*
 457 *Ecol.* 446, 166–176.

458 Merchant, N.D., Fristrup, K.M., Johnson, M.P., Tyack, P.L., Witt, M.J., Blondel, P., Parks, S.E., 2015.
 459 Measuring acoustic habitats. *Methods Ecol. Evol.* 6, 257–265. [https://doi.org/10.1111/2041-](https://doi.org/10.1111/2041-210X.12330)
 460 [210X.12330](https://doi.org/10.1111/2041-210X.12330)

461 Meyer, E., Clapp, M., McKenna, M.F., Crooks, K.R., Buxton, R.T., Angeloni, L.M., Stabenau, E.,
 462 Wittemyer, G., 2018. Efficacy of extracting indices from large-scale acoustic recordings to
 463 monitor biodiversity. *Conserv. Biol.* 32, 1174–1184. <https://doi.org/10.1111/cobi.13119>

464 Miralles, L., Ardura, A., Arias, A., Borrell, Y.J., Clusa, L., Dopico, E., de Rojas, A.H., Lopez, B., Muñoz-
 465 Colmenero, M., Roca, A., Valiente, A.G., Zaiko, A., Garcia-Vazquez, E., 2016. Barcodes of marine
 466 invertebrates from north Iberian ports: Native diversity and resistance to biological invasions.
 467 *Mar. Pollut. Bull.* 112, 183–188. <https://doi.org/10.1016/j.marpolbul.2016.08.022>

468 Nedelec, S.L., Simpson, S.D., Holderied, M.W., Radford, A.N., Lecellier, G., Radford, C.A., Lecchini, D.,
 469 2015. Soundscapes and living communities in coral reefs: temporal and spatial variation. *Mar.*
 470 *Ecol. Prog. Ser.* 524, 125–135. <https://doi.org/10.3354/meps11175>

471 Picciulin, M., Bolgan, M., Codarin, A., Fiorin, R., Zucchetta, M., Malavasi, S., 2013. Passive acoustic
 472 monitoring of *Sciaena umbra* on rocky habitats in the Venetian littoral zone. *Fish. Res.* 145, 76–

473 81. <https://doi.org/10.1016/j.fishres.2013.02.008>

474 Pieretti, N., Duarte, M.H.L., Sousa-Lima, R.S., Rodrigues, M., Farina, A., 2015. Determining temporal
 475 sampling schemes for passive acoustic studies in different tropical ecosystems. *Trop. Conserv.*
 476 *Sci.* 88, 215–234.

477 Pieretti, N., Farina, A., 2013. Application of a recently introduced index for acoustic complexity to an
 478 avian soundscape with traffic noise. *J. Acoust. Soc. Am.* 134, 891–900.
 479 <https://doi.org/10.1121/1.4807812>

480 Pieretti, N., Farina, A., Morri, D., 2011. A new methodology to infer the singing activity of an avian
 481 community: The Acoustic Complexity Index (ACI). *Ecol. Indic.* 11, 868–873.
 482 <https://doi.org/10.1016/j.ecolind.2010.11.005>

483 Pieretti, N., Lo Martire, M., Farina, A., Danovaro, R., 2017. Marine soundscape as an additional
 484 biodiversity monitoring tool: A case study from the Adriatic Sea (Mediterranean Sea). *Ecol.*
 485 *Indic.* 83, 13–20. <https://doi.org/10.1016/j.ecolind.2017.07.011>

486 Pijanowski, B.C., Villanueva-Rivera, L.J., Dumyahn, S.L., Farina, A., Krause, B.L., Napoletano, B.M.,
 487 Gage, S.H., Pieretti, N., 2011. Soundscape Ecology: The Science of Sound in the Landscape.
 488 *Bioscience* 61.

489 Radford, A.N., Kerridge, E., Simpson, S.D., 2014. Behavioral Ecology Acoustic communication in a
 490 noisy world: can fish compete with anthropogenic noise? *Behav. Ecol.* 25, 1022–1030.
 491 <https://doi.org/10.1093/beheco/aru029>

492 Radford, C.A., Jevs, A.G., Tindle, C.T., Montgomery, J.C., 2008. Temporal patterns in ambient noise of
 493 biological origin from a shallow water temperate reef. *Oecologia* 156, 921–929.
 494 <https://doi.org/10.1007/s00442-008-1041-y>

495 Radford, C.A., Stanley, J.A., Simpson, S.D., Jeffs, A.G., 2011. Juvenile coral reef fish use sound to
 496 locate habitats. *Coral Reefs* 30, 295–305.

497 Ricci, S.W., Eggleston, D.B., Bohnenstiehl, D.R., 2017. Use of passive acoustic monitoring to
 498 characterize fish spawning behavior and habitat use within a complex mosaic of estuarine
 499 habitats. *Bull. Mar. Sci.* 93, 439–453. <https://doi.org/10.5343/bms.2016.1037>
 500 Rivers, J.A., 1997. Blue whale, *Balaenoptera musculus*, vocalizations from the waters off central
 501 California, *Marine Mammal Science*.
 502 Rombouts, I., Simon, N., Aubert, A., Cariou, T., Feunteun, E., Guérin, L., Hoebeke, M., McQuatters-
 503 Gollop, A., Rigaut-Jalabert, F., Artigas, L.F., 2019. Changes in marine phytoplankton diversity:
 504 Assessment under the Marine Strategy Framework Directive. *Ecol. Indic.* 102, 265–277.
 505 <https://doi.org/10.1016/j.ecolind.2019.02.009>
 506 Rowley, S.J., 2008. *Inachus phalangium* Leach's spider crab. [WWW Document]. *Mar. Life Inf. Netw.*
 507 *Biol. Sensit. Key Inf. Rev.*
 508 Samaran, F., Stafford, K.M., Branch, T.A., Gedamke, J., Royer, J.-Y., Dziak, R.P., Guinet, C., 2013.
 509 Seasonal and Geographic Variation of Southern Blue Whale Subspecies in the Indian Ocean.
 510 *PLoS One* 8. <https://doi.org/10.1371/journal.pone.0071561>
 511 Sheehan, E. V., Cousens, S.L., Nancollas, S.J., Stauss, C., Royle, J., Attrill, M.J., 2013a. Drawing lines at
 512 the sand: Evidence for functional vs. visual reef boundaries in temperate Marine Protected
 513 Areas. *Mar. Pollut. Bull.* 76, 194–202. <https://doi.org/10.1016/j.marpolbul.2013.09.004>
 514 Sheehan, E. V., Stevens, T.F., Attrill, M.J., 2010. A Quantitative, Non-Destructive Methodology for
 515 Habitat Characterisation and Benthic Monitoring at Offshore Renewable Energy Developments.
 516 *PLoS One* 5. <https://doi.org/10.1371/journal.pone.0014461>
 517 Sheehan, E. V., Stevens, T.F., Gall, S.C., Cousens, S.L., Attrill, M.J., Fulton, C.J., 2013b. Recovery of a
 518 Temperate Reef Assemblage in a Marine Protected Area following the Exclusion of Towed
 519 Demersal Fishing. *PLoS One* 8, e83883. <https://doi.org/10.1371/journal.pone.0083883>
 520 Staatterman, E., Ogburn, M.B., Altieri, A.H., Brandl, S.J., Whippo, R., Seemann, J., Goodison, M., Duffy,

521 J.E., 2017. Bioacoustic measurements complement visual biodiversity surveys: preliminary
 522 evidence from four shallow marine habitats. *Mar. Ecol. Prog. Ser.* 575, 207–215.
 523 <https://doi.org/10.3354/meps12188>

524 Staaterman, E., Paris, C.B., DeFerrari, H.A., Mann, D.A., Rice, A.N., D'Alessandro, E.K., 2014. Celestial
 525 patterns in marine soundscapes. *Mar. Ecol. Prog. Ser.* 508, 17–32.
 526 <https://doi.org/10.3354/meps10911>

527 Stevens, T.F., Sheehan, E. V., Gall, S.C., Fowell, S.C., Attrill, M.J., 2014. Monitoring benthic
 528 biodiversity restoration in Lyme Bay marine protected area: Design, sampling and analysis.
 529 *Mar. Policy* 45, 310–317. <https://doi.org/10.1016/j.marpol.2013.09.006>

530 Sueur, J., Aubin, T., Simonis, C., 2008a. SEEWAVE, A FREE MODULAR TOOL FOR SOUND ANALYSIS
 531 AND SYNTHESIS EQUIPMENT. *Bioacoustics* 18, 213–226.
 532 <https://doi.org/10.1080/09524622.2008.9753600>

533 Sueur, J., Farina, A., 2015. Ecoacoustics: the Ecological Investigation and Interpretation of
 534 Environmental Sound. *Biosemiotics* 8, 493–502. <https://doi.org/10.1007/s12304-015-9248-x>

535 Sueur, J., Pavoine, S., Hamerlynck, O., Duvail, S., 2008b. Rapid acoustic survey for biodiversity
 536 appraisal. *PLoS One* 3. <https://doi.org/10.1371/journal.pone.0004065>

537 Trenkel, V.M., Ressler, P.H., Jech, M., Giannoulaki, M., Taylor, C., 2011. Underwater acoustics for
 538 ecosystem-based management: state of the science and proposals for ecosystem indicators.
 539 *Mar. Ecol. Prog. Ser.* 442, 285–301. <https://doi.org/10.3354/meps09425>

540 Vermeij, M.J.A., Marhaver, K.L., Huijbers, C.M., Nagelkerken, I., Simpson, S.D., 2010. Coral Larvae
 541 Move toward Reef Sounds. *PLoS One* 5, e10660. <https://doi.org/10.1371/>

542 Wabnitz, C.C.C., Lam, V.W.Y., Reygondeau, G., Teh, L.C.L., Al-Abdulrazzak, D., Khalfallah, M., Pauly,
 543 D., Deng Palomares, M.L., Zeller, D.C., Cheung, W.W.L., 2018. Climate change impacts on
 544 marine biodiversity, fisheries and society in the Arabian Gulf. *PLoS One* 13, 1–26.

545 <https://doi.org/10.1371/journal.pone.0194537>

546 Worm, B., Barbier, E.B., Beaumont, N., Duffy, J.E., Folke, C., Halpern, B.S., Jackson, J.B.C., Lotze, H.K.,
547 Micheli, F., Palumbi, S.R., Sala, E., Selkoe, K.A., Stachowicz, J.J., Watson, R., 2006. Impacts of
548 Biodiversity Loss on Ocean Ecosystem Services. *Science* (80-.). 314, 787–790.

549